


The effects of age on dyspnea and respiratory mechanical and neural responses to exercise in healthy men

William MacAskill^{1,2,3,4}  | Ben Hoffman^{1,3} | Michael A. Johnson⁵ |
 Graham R. Sharpe⁵ | Joshua Rands^{1,2} | Shoena E. Wotherspoon⁶ |
 Yaroslav Gevorkov⁷ | Tracy L. Kolbe-Alexander^{1,3,8} | Dean E. Mills^{1,2,3}

¹School of Health and Medical Sciences, University of Southern Queensland, Ipswich, Queensland, Australia

²Respiratory and Exercise Physiology Research Group, School of Health and Wellbeing, University of Southern Queensland, Ipswich, Queensland, Australia

³Centre for Health Research, Institute for Resilient Regions, University of Southern Queensland, Ipswich, Queensland, Australia

⁴Rural Clinical School, Griffith University, Toowoomba, Queensland, Australia

⁵Exercise and Health Research Group, Sport, Health and Performance Enhancement (SHAPE) Research Centre, School of Science and Technology, Nottingham Trent University, Nottinghamshire, UK

⁶Darling Downs Health, Queensland Health, Queensland, Australia

⁷Institute of Vision Systems, Hamburg University of Technology, Hamburg, Germany

⁸UCT Research Centre for Health through Physical Activity, Lifestyle and Sport (HPALS), Division of Research Unit for Exercise Science and Sports Medicine, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

Correspondence

Dean E. Mills, School of Health and Medical Sciences, University of Southern Queensland, Ipswich, Queensland, Australia.

Email: dean.mills@usq.edu.au

Funding information

Australian Government, Research Training Program

Abstract

The respiratory muscle pressure generation and inspiratory and expiratory neuromuscular recruitment patterns in younger and older men were compared during exercise, alongside descriptors of dyspnea. Healthy younger ($n = 8$, 28 ± 5 years) and older ($n = 8$, 68 ± 4 years) men completed a maximal incremental cycling test. Esophageal, gastric (P_{ga}) and transdiaphragmatic pressures, and electromyography (EMG) of the crural diaphragm were measured using a micro-transducer and EMG catheter. EMG of the parasternal intercostals, sternocleidomastoids, and rectus abdominis were measured using skin surface electrodes. After the exercise test, participants completed a questionnaire to evaluate descriptors of dyspnea. P_{ga} at end-expiration, P_{ga} expiratory tidal swings, and the gastric pressure-time product (PTP_{ga}) at absolute and relative minute ventilation were higher ($p < 0.05$) for older compared to younger men. There were no differences in EMG responses between older and younger men. Younger men were more likely to report shallow breathing ($p = 0.005$) than older men. Our findings showed younger and older men had similar respiratory neuromuscular activation patterns and reported different dyspnea descriptors, and that older men had greater expiratory muscle pressure generation during exercise. Greater expiratory muscle pressures in older

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Physiological Reports* published by Wiley Periodicals LLC on behalf of The Physiological Society and the American Physiological Society.

men may be due to compensatory mechanisms designed to offset increasing airway resistance due to aging. These results may have implications for exercise-induced expiratory muscle fatigue in older men.

KEYWORDS

aging, dyspnea, exercise, mechanical, neural

1 | INTRODUCTION

Healthy aging results in progressive changes to the lungs, airways, chest wall, and respiratory muscles, leading to a decline in respiratory structure and function (Janssens et al., 1999; Lalley, 2013; Meiners et al., 2015; Roman et al., 2016). The changes that occur during healthy aging include reductions in chest wall compliance due to structural changes in the thoracic cage (Estenne et al., 1985; Mittman et al., 1965) including kyphosis, calcification of the costal cartilage, narrowing of intervertebral disc spaces, and osteoporosis (Holcombe et al., 2017). Additionally, lung compliance progressively increases (Turner et al., 1968), possibly due to deterioration in the spatial arrangement and/or cross linking of the elastic fiber network (Fukuchi, 2009). Respiratory muscle function also declines with age, due to the atrophy and denervation of type II muscle fibers (Elliott et al., 2016; Ticinesi et al., 2017). There is evidence that this also occurs within the diaphragm due to a loss of larger phrenic motoneurons (Fogarty et al., 2018; Prakash & Sieck, 1998).

Inspiratory muscle pressure generation and the resistive and elastic components of the work of breathing (WOB) are greater for older compared to younger adults during maximal incremental exercise (Molgat-Seon et al., 2018b, Molgat-Seon et al., 2019, Weavil et al., 2022, Molgat-Seon et al., 2018a). Molgat-Seon et al. (2018b) reported that inspiratory neuromuscular recruitment patterns during maximal incremental exercise are different between healthy younger and older adults. Older adults had greater esophageal and transdiaphragmatic pressure-time products (PTP_{es} and PTP_{di}) at absolute minute ventilations (\dot{V}_E) compared to younger adults (Molgat-Seon et al., 2018b). Older adults also had greater recruitment of primary and accessory inspiratory muscles, evidenced by their higher amplitudes for crural diaphragm electromyography (EMG_{di}), scalene EMG and sternocleidomastoids EMG (EMG_{scm}) at absolute \dot{V}_E (Molgat-Seon et al., 2018b). At relative \dot{V}_E , EMG_{scm} was also higher for older compared to younger adults (Molgat-Seon et al., 2018b). While the work of Molgat-Seon et al. (2018b) provides valuable insight into the differences between younger and older adults in inspiratory muscle pressure generation and neuromuscular recruitment patterns during exercise, they did

not report on the contributions of the expiratory muscles. With advancing age, the muscle mass of the rectus abdominis declines (Abe et al., 2011; Kubo, 1994) and abdominal wall compliance is reduced (Estenne et al., 1985). Therefore, compared to younger men, greater abdominal muscle recruitment may be expected in older men to support the inspiratory role of the abdominal musculature, and to compensate for their greater expiratory flow limitation and potentially higher expiratory WOB during exercise (Smith et al., 2018; Weavil et al., 2022).

Dyspnea is defined as “a subjective experience of breathing discomfort that consists of qualitatively distinct sensations that vary in intensity” (American Thoracic Society, 1999, Parshall et al., 2012). Higher ratings of perceived exertion (RPE) for dyspnea have been reported in older compared to younger men (Ofir et al., 2008), although others report no difference (Faisal et al., 2015). Dyspnea is influenced by a wide variety of stimuli and sensory receptors which contributes to the challenges in fully understanding its mechanistic causes (Laviolette & Laveneziana, 2014; Parshall et al., 2012). Ofir et al. (2008) compared descriptors of dyspnea between younger and older men and women following maximal incremental treadmill exercise and found that shallow breathing was reported by a greater proportion of older men and women. Breathing discomfort was also higher for older than younger women. Older men only reported greater breathing discomfort than young men at higher relative oxygen uptake (≥ 25 mL/kg/min) (Ofir et al., 2008). Descriptors of dyspnea may provide a more complete understanding of its subjective experience and may assist in developing a more comprehensive understanding of dyspnea (Lansing et al., 2009). Due to age-related changes in respiratory physiology, older men may report a different affective experience of dyspnea during exercise. No published literature has measured detailed respiratory muscle pressure generation and neuromuscular recruitment patterns, including expiratory measurements, alongside descriptors of dyspnea in younger and older men during incremental exercise.

The aim of this study was to compare respiratory muscle pressure generation, inspiratory and expiratory neuromuscular recruitment patterns, and descriptors of dyspnea, in younger and older men during exercise. We

hypothesized that older men would have greater expiratory muscle recruitment and expiratory muscle pressure generation during exercise and select descriptors of dyspnea consistent with an increased respiratory muscle pressure generation following exercise.

2 | METHODS

2.1 | Participants

Healthy younger ($n=8$) and older ($n=8$) men participated in this study (Table 1). The exclusion criteria were current cigarette smokers (two older adults quit smoking more than 10 years ago); history or current symptoms of cardiopulmonary disease; prescribed medications affecting the cardiopulmonary system; contraindications to exercise testing; and a body mass index of below 18.5 or above 30 kg/m². Participants were aged between 18–35 (younger) or 65–80 (older) years and recreationally active. The definition of a recreationally active participant for this study was participation in nonprofessional sport one to three times per week. The study was approved by the University of Southern Queensland's Human Research Ethics Committee and all procedures conformed to the standards set by the Declaration of Helsinki, except for

TABLE 1 Participant characteristics and pulmonary function for the younger and older men.

	Younger men	Older men	p-value
Age, years	28 ± 5	68 ± 4	<0.001
Height, cm	179 ± 6	180 ± 7	0.853
Body mass, kg	83 ± 9	86 ± 14	0.707
BMI, kg/m ²	26 ± 3	26 ± 3	0.787
FVC, L	5.2 ± 0.8	4.5 ± 1.1	0.153
FVC, %predicted	98 ± 12	102 ± 16	0.589
FEV ₁ , L	4.1 ± 0.6	3.3 ± 0.5	0.007
FEV ₁ , %predicted	95 ± 13	99 ± 14	0.579
FEV ₁ /FVC, %	81 ± 6	75 ± 10	0.136
FEV ₁ /FVC, %predicted	97 ± 8	99 ± 13	0.856
P _{es,max} , cmH ₂ O	−131 ± 52	−112 ± 39	0.432
P _{ga,max} , cmH ₂ O	138 ± 71	132 ± 45	0.853
P _{di,max} , cmH ₂ O	71 ± 7	79 ± 11	0.096

Values are mean ± SD.

Abbreviations: BMI, body mass index; FVC, forced vital capacity; FEV₁, forced expiratory volume in 1 s; P_{es,max}, maximum esophageal pressure; P_{ga,max}, maximum gastric pressure; P_{di,max}, maximum transdiaphragmatic pressure. P_{es,max} and P_{di,max} were collected during a maximal inspiratory pressure maneuver. P_{ga,max} was collected during a maximal expiratory pressure maneuver.

registration in a database. The trial was registered with the Australian and New Zealand Clinical Trial Registry (ACTRN 376141). All participants provided written, informed consent prior to participation.

2.2 | Experimental design

The study utilized a cross-sectional design. Each participant visited the laboratory on two occasions, separated by a minimum of 48 h and a maximum of 1 week. Visit 1 included screening for the eligibility criteria, anthropometric and pulmonary function assessment, and familiarization with all the experimental procedures. Visit 2 included a maximal incremental cycling test with measurements of respiratory muscle pressures and EMG, and ventilatory, cardiovascular and perceptual responses to exercise. Participants were instructed to abstain from food (4 h), caffeine (12 h), and strenuous exercise (48 h) before testing.

2.3 | Anthropometrical measures and pulmonary function

Height and body mass were recorded using a wall mounted electronic stadiometer (Seca 213; Seca) and an electronic scale (Tanita BC-541), respectively. Body mass index was then calculated as body mass divided by height in meters squared. Pulmonary function was assessed using a spirometer (Vmax[®] Encore PFT system; Vyaire Medical) according to published guidelines (Miller et al., 2005). Pulmonary function measurements were expressed as absolute values and as percentages of predicted values (Quanjer et al., 2012).

2.4 | Maximal incremental cycling

Maximal incremental cycling was conducted on an electronically braked cycle ergometer (Corival CPET; Lode) automatically controlled by software (SentrySuite; Vyaire Medical). The protocol consisted of a baseline resting period of 5 min (while participants were seated on the ergometer) followed by 1 min of unloaded cycling. The initial work rate was individualized for each participant, with participants beginning at either 40, 60, or 80 W, depending upon their age, and self-reported exercise training history. Work rate was increased by 20 W every 2 min. Participants maintained a constant self-selected cadence above 60 revs/min. Exercise ceased at the limit of tolerance or when cycling cadence could not be maintained above 60 revs/min.

2.5 | Respiratory pressures and electromyography

Esophageal pressure (P_{es}), gastric pressure (P_{ga}), transdiaphragmatic pressure (P_{di}), and EMG_{di} were measured in real time using a micro-transducer and EMG esophageal catheter as previously described (MacAskill et al., 2021). The catheter housed two pressure transducers ($\sim 5 \times 2$ mm), separated by 22.8 cm, which were constructed using half bridge thin film resistive strain gauge sensors coated with a silicone elastomer with frequency responses of 10–20 kHz. The catheter comprised a 100 cm silicon shaft (2.7 mm diameter) containing nine silver electrodes spaced 1 mm apart and the pressure transducers were positioned proximally and distally to the electrodes. The nine electrodes formed five pairs (i.e., electrodes 1 & 5; 2 & 6; 3 & 7; 4 & 8; 5 & 9), with each pair recording a difference in electrical potential across the crural diaphragm. The highest mean EMG_{di} recorded from any of the catheters five EMG pairs was utilized for analysis (Dacha et al., 2019).

Prior to instrumentation, the catheter was soaked in water for 1 h as per manufacturer's instructions to reduce baseline drift. The micro-transducer catheter was then placed inside a small section of airtight plastic tubing and calibrated by injecting or withdrawing air, via a three-way open valve connected to a glass syringe and a handheld respiratory pressure meter (Micro RPM; Vyair Medical). P_{es} and P_{ga} were calibrated from -100 cmH₂O to $+100$ cmH₂O. Catheter placement was preceded by the administration of 91% aerosolized lidocaine hydrochloride and 9% phenylephrine hydrochloride (Co-Phenylcaine Forte Spray; Almed). The catheter was passed peri-nasally into the stomach until a negative deflection in P_{es} and a positive deflection in P_{ga} were observed during repeated sniffs. Further sniffs were performed (to monitor P_{es} and P_{ga} inflection) while the catheter position was optimized to record maximal EMG_{di} on the superior and inferior electrode pairs. An occlusion test was then performed to confirm the catheters location in the esophagus before securing the catheter in position (Baydur et al., 1982).

P_{di} was calculated by subtracting P_{es} from P_{ga} . P_{es} , P_{ga} and P_{di} were recorded throughout exercise. Additionally, these signals were integrated for active pressures over each breath's periods of inspiration (P_{di} and P_{es}) and expiration (P_{ga}), then multiplied by breathing frequency to calculate PTP_{di} , PTP_{es} , and gastric pressure time products (PTP_{ga}). Pressures were normalized using the maximum pressure obtained from any forced vital (for P_{ga}) or inspiratory (for P_{di} and P_{es}) capacity maneuvers performed during pulmonary function assessment or exercise testing.

Skin-surface parasternal EMG (EMG_{para}), EMG_{scm} , and EMG rectus abdominis (EMG_{ra}) were assessed using pairs of bipolar skin-surface electrodes (Ambu WhiteSensor

40,713). The electrodes were placed on the right side of the torso after shaving (if required), cleaning and abrading the skin with skin preparation gel (NuPrep; Weaver and Company) and alcohol wipes. The electrodes were 2 cm in diameter and an interelectrode distance of 2 cm was utilized. Surface electrodes for EMG_{para} and EMG_{scm} were placed as described previously (Ramsook et al., 2016). EMG_{para} were placed in the space of the second rib roughly 3 cm lateral to the sternum, and for EMG_{scm} at the midpoint along the longitudinal axis of the sternocleidomastoid muscle between the mastoid process and the medial clavicle. EMG_{ra} were placed 2 cm superior and 2–4 cm lateral to the umbilicus (Fuller et al., 1996; Ng et al., 1998). Optimal electrode positioning was determined by utilizing maximal inspiratory (EMG_{para} and EMG_{scm}) and expiratory (EMG_{ra}) mouth pressure measurements to evoke EMG responses. EMG data were transformed into root mean square (RMS) with a time constant of 0.05 s. EMG RMS values were manually selected between the QRS complexes of the electrocardiogram (ECG) to limit the effect of cardiac artifacts on respiratory muscle recruitment (Guenette et al., 2014; Jolley et al., 2015; Schaeffer et al., 2014). EMG was normalized using the maximum EMG obtained from any forced vital (EMG_{ra}) or inspiratory (EMG_{di} , EMG_{para} and EMG_{scm}) capacity maneuvers performed during pulmonary function assessment or exercise testing.

2.6 | Metabolic, Ventilatory, and cardiorespiratory measurements

Participants wore a facemask (Model 7940; Hans Rudolph) which was tightly fitted to minimize leaks and connected to a turbine flow sensor (Digital volume transducer; Vyair Medical) that was calibrated using a 3 L syringe. Pulmonary gas exchange was measured breath by breath using a metabolic cart (Vmax[®] Encore PFT system; Vyair Medical). As the metabolic cart could not be integrated with the data acquisition unit, flow measurements were additionally recorded with a heated calibrated pneumotachograph (Model 3813; Hans Rudolph) and were time aligned with pressure and EMG data. The pneumotachograph was attached distally to the flow sensor from the metabolic cart. This assembly (facemask, pneumotach, and flow sensor) has an estimated total dead space of 144 mL. Cardiac frequency and ECG responses were monitored continuously during testing using a 12-lead ECG (PC-ECG 1200; Norav). Arterial oxygen saturation was estimated using infrared fingertip pulse oximetry (Model 8600; Nonin). The maximal work rate was calculated as the work rate of the penultimate stage plus the fraction of work completed in the final stage. The highest oxygen

uptake recorded during any 30s period defined maximal oxygen uptake ($\dot{V}O_{2\max}$). Predicted $\dot{V}O_{2\max}$ was determined using a reference equation for healthy men undertaking cycling exercise (de Souza e Silva et al., 2018). Predicted maximal cardiac frequency was determined using published formulas for healthy men free of cardiovascular disease (Ozemek et al., 2017).

2.7 | Operational lung volumes

Volume was obtained by numerical integration of the flow signal. Operational lung volumes were quantified by measuring inspiratory capacity (IC) relative to forced vital capacity (FVC). During the resting stage, participants performed FVC and IC maneuvers in duplicate. Participants performed further IC maneuvers in duplicate during the exercise testing within the final 1 min of each incremental stage. Strong verbal encouragement was given during each maximal inspiratory effort maneuver.

2.8 | Perceptual responses

Intensities of breathing (dyspnea) and leg discomfort were rated using Borg's CR-10 RPE scale (Borg, 1982). Breathing discomfort was defined as "a feeling of labored or difficult breathing," and leg discomfort as "a feeling of fatigue in the leg muscles." In this scale 0 represents "no discomfort" while 10 represents "maximum imaginable discomfort" (Borg, 1982). Reporting was completed at rest, at the end of each stage, and immediately following the cessation of exercise. After the exercise test, participants completed a questionnaire to evaluate descriptors of dyspnea. This questionnaire was a modified version of those previously utilized by Simon et al. (1990) and Cory et al. (2015). Participants were asked to indicate whether each descriptor was applicable or not and their responses were recorded.

2.9 | Data capture and analysis of exercise end points

Raw pressure and EMG data were amplified with a Quad Bridge Amp FE224 and Octal Bio Amp FE238 (AD Instruments), respectively. Pressure data were low-pass filtered at 1 kHz and EMG data were high-pass filtered at 80 Hz. Pressure, EMG, and flow data were sampled continuously at 10 kHz using a 16-channel analog-to-digital data acquisition system (PowerLab 16/35; AD Instruments) and recorded using LabChart v8.1.2 software (AD Instruments).

Nonphysiological pressure, EMG, and flow data that resulted from swallowing, coughing, and breath holding were identified by visual inspection and removed. All physiological variables collected during the cycle ergometer test were averaged in 30s epochs. Data were collected in the final 1 min of each 2 min stage and used for statistical analysis. During this time, participants were asked to look forward, minimize any head or neck movement, keep a loose grip on the handlebars, and to avoid talking or swallowing. Five representative and consecutive breaths were selected from the final 30s of each 2 min stage for analysis (Dacha et al., 2019).

The schedule of data collection in each 2 min exercise stage was as follows: 0–60s, exercise only; 61–90s, IC maneuvers; and 91–120s, once breathing pattern returned to baseline perceptual responses and representative breaths were collected.

2.10 | Statistical analyses

Statistical analyses were performed using SPSS for Windows (IBM). An initial power calculation was performed based on differences in P_{ga} swings during exercise between younger and older men. Power analysis indicated that a sample size of 10 (5 younger and 5 older men) would be required to detect differences in P_{ga} swings ($\alpha=0.05$, $\text{power}=0.8$) and provide a large effect size ($d_z=2.1$).

Normality was assessed using a Shapiro–Wilk test. Comparisons between younger and older men for characteristics, pulmonary function and peak metabolic, ventilatory and cardiorespiratory responses were determined using an independent *t*-test and a Mann–Whitney *U*-test for parametric and nonparametric data, respectively. The older and younger men were compared at rest and at matched \dot{V}_E of ~40, 50, 60, and 70 L/min (31–45, 46–55, 56–65, 66–75 L/min), as 70 L/min was the highest \dot{V}_E reached by all participants. The older and younger men were also compared at matched relative \dot{V}_E of ~20, 40, 60, 80 and 100% of maximum (11–30, 31–50, 51–70, 71–99, 100% maximum). Interpolation was used if necessary to match participant \dot{V}_E .

Between-group differences were analyzed with a two-way analysis of variance to determine the effects of "age" (younger vs. older) and " \dot{V}_E " (absolute \dot{V}_E : rest and at ~40, 50, 60, and 70 L/min; or relative \dot{V}_E : ~20, 40, 60, 80, and 100% maximum). Significant main effects of age and age \times ventilation interaction effects were followed by planned pairwise comparisons between ages using the Bonferroni post hoc analysis. Comparisons between younger and older participants for categorical descriptors of dyspnea were made using Fisher's Exact test. Statistical

significance was set at $p \leq 0.05$. Values are expressed as means \pm SD.

3 | RESULTS

3.1 | Participant characteristics and pulmonary function

There were no differences in participant characteristics and pulmonary function between younger and older men, apart from forced expiratory volume in 1 s (FEV_1) which was lower for older men (Table 1).

3.2 | Metabolic, Ventilatory, and cardiorespiratory responses

The metabolic, ventilatory and cardiorespiratory responses for the younger and older men at peak exercise are shown in Table 2. At peak exercise, the percentage of predicted $\dot{V}O_{2max}$ was higher, whereas carbon dioxide output, respiratory exchange ratio, minute ventilation, $\dot{V}_E/\dot{V}O_2$ and cardiac frequency were lower, in older compared to younger men. Significant age \times ventilation interaction effects were not observed at either absolute or relative \dot{V}_E .

3.3 | Respiratory pressure generation

Absolute P_{es} , P_{ga} , and P_{di} and pressure swings are shown in Figure 1. Pressures as a percentage of maximum are shown in Figure 2. PTPs are shown in Figure 3. Main effects of age and age \times ventilation interaction effects are shown in the figures. All measurements increased during exercise (main effect of absolute and relative \dot{V}_E , $p < 0.01$), except for P_{di} at end-expiration (absolute \dot{V}_E , $P = 0.431$; relative \dot{V}_E , $p = 0.354$).

Main effects of age were observed for P_{ga} at end-expiration and P_{ga} swing (at absolute and relative \dot{V}_E) and PTP_{ga} (at relative \dot{V}_E). This effect showed that older men's P_{ga} at end-expiration was higher, and P_{ga} swings larger (Figure 1). Age \times ventilation interactions were observed for P_{ga} at end-expiration and PTP_{ga} (at relative and absolute \dot{V}_E) and for P_{ga} swing (at relative \dot{V}_E). Pairwise differences in P_{ga} at end-expiration and P_{ga} swings were apparent from $\dot{V}_E = 50$ L/min and $\dot{V}_E = 60\%$ until the end of the exercise test ($p < 0.05$). Pairwise differences in PTP_{ga} occurred from $\dot{V}_E = 80\%$ until the end of the exercise test ($p < 0.05$). When P_{ga} tidal swings were expressed as a percentage of maximum, an age \times ventilation effect was observed (at absolute and relative \dot{V}_E), indicating greater swings in older men (Figure 2). Age \times ventilation

TABLE 2 Metabolic, ventilatory, and cardiorespiratory responses for younger and older men at peak exercise.

	Younger men	Older men	p-value
W_{peak} , W	212 \pm 52	206 \pm 37	0.816
$\dot{V}O_2$, L/min	3.22 \pm 0.64	2.55 \pm 1.15	0.174
$\dot{V}O_2$, mL/kg/min	37.3 \pm 5.6	33.7 \pm 8.5	0.330
$\dot{V}O_2$, %predicted	87.8 \pm 17.3	115.2 \pm 26.8	0.030
$\dot{V}CO_2$, L/min	3.56 \pm 0.70	2.51 \pm 1.05	0.033
RER	1.18 \pm 0.08	1.02 \pm 0.16	0.024
PET O_2 , mmHg	119 \pm 5	117 \pm 7	0.355
PET CO_2 , mmHg	33.7 \pm 3.4	31.6 \pm 6.2	0.408
\dot{V}_E , L/min	130 \pm 24	94 \pm 37	0.034
$\dot{V}_E/\dot{V}O_2$	41.3 \pm 1.6	37.0 \pm 4.6	0.047
$\dot{V}_E/\dot{V}CO_2$	34.7 \pm 2.7	38.2 \pm 7.3	0.312
F_b , breaths/min	50 \pm 10	38 \pm 13	0.119
V_T , L	2.76 \pm 0.55	2.57 \pm 0.80	0.594
V_T , % FVC	53 \pm 7	64 \pm 13	0.065
T_I , s	0.61 \pm 0.11	0.69 \pm 0.11	0.180
T_{TOT} , s	1.24 \pm 0.22	1.46 \pm 0.23	0.072
T_I/T_{TOT}	0.48 \pm 0.01	0.47 \pm 0.02	0.656
IC, L	3.45 \pm 0.82	3.24 \pm 0.82	0.660
F_c , beats/min	179 \pm 14	149 \pm 13	0.001
F_c , %predicted	92 \pm 7	93 \pm 8	0.098
Sp O_2 , %	95 \pm 2	95 \pm 2	0.611

Values are mean \pm SD.

Abbreviations: W_{peak} , peak work rate; $\dot{V}O_2$, oxygen uptake; $\dot{V}CO_2$, carbon dioxide output; RER, respiratory exchange ratio; PET O_2 , end tidal oxygen pressure; PET CO_2 , end tidal carbon dioxide pressure; \dot{V}_E , minute ventilation; F_b , breathing frequency; V_T , tidal volume; T_I , inspiratory time, T_{TOT} , total breath time; T_I/T_{TOT} , duty cycle; IC, inspiratory capacity; FVC, forced vital capacity; F_c , cardiac frequency; Sp O_2 , estimated arterial oxygen saturation.

interactions were also observed at absolute \dot{V}_E for P_{es} at end-expiration, P_{di} swing, and P_{di} swing as a percentage of maximum.

3.4 | Respiratory muscle electromyography

EMG responses at rest and during exercise for the younger and older men are shown in Figure 4, along with main effects of age and age \times ventilation interaction effects. EMG $_{di}$, EMG $_{scm}$, EMG $_{para}$ and EMG $_{ra}$ measurements increased during exercise (main effects of absolute and relative \dot{V}_E , $p < 0.01$). No main effects of age or age \times ventilation interaction effects were observed, indicating neuromuscular activation patterns in response to exercise were similar in both age groups.




FIGURE 1 Esophageal (P_{es} ; a), gastric (P_{ga} ; b) and transdiaphragmatic (P_{di} ; c) pressures at rest and during incremental exercise. End-inspiration (P_{insp} , solid lines) and expiration (P_{exp} , dashed lines) are shown for absolute (left hand panels) and relative (right hand panels) minute ventilation for younger (open circles) and older men (filled boxes). Tidal pressure swings (P_{swing}) are represented by the pressure difference between points of end-inspiration and end-expiration and are shown for younger (blue fill) and older (diagonal fill) men. Data are presented as mean \pm SD. Main effects of age and age \times ventilation interaction effects are provided in each panel. Significant difference in pressures at (*) end-expiration and (†) tidal pressure swings between younger and older men ($p < 0.05$).

3.5 | Perceptual responses

RPE for dyspnea and leg discomfort at rest and during exercise are shown in Figure 5 with main effects of age and age \times ventilation interaction effects. RPE for dyspnea and leg discomfort increased during exercise (main effects of

absolute and relative \dot{V}_E , $p < 0.001$). There were no differences between older and younger men for leg discomfort at absolute and relative \dot{V}_E . Older men reported greater dyspnea than younger men during exercise at absolute \dot{V}_E , but no differences were present for dyspnea for relative \dot{V}_E .




FIGURE 2 Esophageal (P_{es} ; a), gastric (P_{ga} ; b), and transdiaphragmatic (P_{di} ; c) pressure swings as a percentage of maximum at rest and during maximal incremental exercise. Relative pressures are expressed relative to maximal P_{ga} . Responses at absolute (left hand panels) and relative (right hand panels) minute ventilation are shown for younger (open circles) and older men (filled boxes). Data are presented as mean \pm SD. Main effects of age and age \times ventilation interaction effects are provided in each panel.

Subjective descriptors at the cessation of exercise for the younger and older men are shown in Figure 6. Compared to older men, a greater proportion of younger men reported that “My breathing feels shallow” ($n=5$ vs. $n=0$; $p=0.021$) and “Breathing out requires more effort” ($n=5$ vs. $n=0$; $p=0.021$). When results were assessed as descriptor clusters, younger men were more likely to report shallow breathing ($n=6$ vs. $n=0$; $p=0.005$) than older men.

4 | DISCUSSION

4.1 | Main findings

The aim of this study was to compare respiratory muscle pressure generation and inspiratory and expiratory neuromuscular recruitment patterns between younger and older men during exercise, alongside descriptors of




FIGURE 3 Esophageal (PTP_{es}; a), gastric (PTP_{ga}; b), and transdiaphragmatic (PTP_{di}; c) pressure–time products at rest and during maximal incremental exercise. Responses at absolute (left hand panels) and relative (right hand panels) minute ventilation are shown for younger (open circles) and older men (filled boxes). Data are presented as mean \pm SD. Main effects of age and age \times ventilation interaction effects are provided in each panel. *Significant difference between younger and older men ($p < 0.05$).

dyspnea. We hypothesized that older men would have greater expiratory muscle recruitment and, an elevated expiratory muscle pressure generation during exercise and select descriptors of dyspnea consistent with an increased respiratory muscle pressure generation following

exercise. The main findings were twofold. First, older men demonstrated greater abdominal muscle pressure generation during exercise compared to younger men. Second, older men reported expiratory difficulty and shallow breathing less often than younger men.




FIGURE 4 Electromyography (EMG) of the crural diaphragm (EMG_{di}; a), sternocleidomastoids (EMG_{scm}; b), parasternals (EMG_{para}; c), and rectus abdominis (EMG_{ra}; d) at rest and during maximal incremental exercise. Responses at absolute (left hand panels) and relative (right hand panels) minute ventilation are shown for younger (open circles) and older men (filled boxes). Panels b-d have a sample size of $n = 7$ for younger men. Data are presented as mean \pm SD. Main effects of age and age \times ventilation interaction effects are provided in each panel.




FIGURE 5 Rating of perceived exertion (RPE) for dyspnea (A,B) and leg discomfort (C,D) at rest and during maximal incremental exercise. Responses at absolute (A,C) and relative (B,D) minute ventilations are shown for younger (open circles) and older men (filled boxes). Data are presented as means \pm SD. Main effects of age and age \times ventilation interaction effects are provided in each panel.




FIGURE 6 Selection frequency of dyspnea descriptors (a) and dyspnea clusters (b) following maximal incremental exercise. Data are shown for younger (open) and older men (filled). *Significantly different between younger and older men ($p < 0.05$).

4.2 | Respiratory pressure generation

We observed at rest that both younger and older men had similar abdominal muscle pressure generation with no

age-based differences present in P_{ga} or its derived measurements. However, from relatively low intensity exercise (≤ 50 L/min or $\leq 60\%$ \dot{V}_E) until the limit of exercise tolerance, P_{ga} at end-expiration and P_{ga} swing (at absolute and

relative \dot{V}_E), and PTP_{ga} (at relative \dot{V}_E) measurements were higher for older compared to younger men. Smith et al. (2018) reported that the expiratory resistive WOB was greater for older compared to younger men during maximal incremental exercise. Our data shows higher P_{ga} and PTP_{ga} which are indicative of a higher expiratory WOB. Our data suggest that older men may be utilizing the abdominal musculature to support expiration by compensating for age-based increases in airway resistance and decreases in chest wall compliance. However, Campbell diagram analysis or more descriptive measures of mechanical ventilatory work would be required to confirm this observation.

Age \times ventilation interactions were observed for P_{es} at end-expiration, P_{di} swing, and P_{di} swing as a percentage of maximum, with higher values for older men than younger men at absolute \dot{V}_E . Despite this, no differences in PTP_{es} or PTP_{di} were observed in older compared to younger men. This finding is in partial agreement with those of Molgat-Seon et al. (2018b) who reported no differences in PTP_{es} or PTP_{di} at relative \dot{V}_E between younger and older adults, with differences between the ages only observed at absolute \dot{V}_E . The lack of difference in PTP_{es} or PTP_{di} at absolute \dot{V}_E may be due in part to the lower $\dot{V}O_{2max}$ of participants in the current study compared to those in the Molgat-Seon et al. (2018b) study. Individuals with a higher $\dot{V}O_{2max}$ may utilize the diaphragm and accessory muscles to a greater extent than untrained individuals because these muscles are trained (Erail et al., 2022) to support higher ventilatory work, and this difference may be more apparent during aging.

4.3 | Respiratory neuromuscular activation patterns

As ventilatory demand increases during exercise, inspiratory and expiratory muscle recruitment is gradually increased to support the increased F_b (Aliverti et al., 1997; Romer & Polkey, 2008). However, little research has been undertaken to understand the impact of healthy aging on respiratory neuromuscular activation patterns. Abraham et al. (2002) measured EMG_{ra} using fine wire intramuscular EMG during incremental and constant load cycling exercise and showed that younger men activated the rectus abdominis early in exercise and that muscle activation plateaued at approximately 20–40% of peak exercise. We also observed an increase in EMG_{ra} during exercise, though no differences were observed between age groups. Why there were no differences in EMG_{ra} , yet greater expiratory pressure generation in older men is not clear. It is possible that the rectus abdominis may contribute to postural control during exercise; however, studies in young men have shown

that the rectus abdominis is primarily activated for respiration as opposed to postural or locomotive needs (Abraham et al., 2002). Another possibility is differences in older and younger men's mechanical advantage and abdominal musculature neuromuscular activation patterns. As the lung inflates during inspiration from functional residual capacity to total lung capacity, the rectus abdominis and external obliques lengthen by 1–3%, the internal obliques by 15%, and the transversus abdominis by 25% (De Troyer and Boriek, 2011). Furthermore, ventilation induced by inhalation of carbon dioxide enriched gas mixtures in humans activates the transversus abdominis and internal obliques before the rectus abdominis and external obliques (De Troyer and Boriek, 2011). Thus, increases in P_{ga} may be predominantly generated by the transversus abdominis and/or the internal obliques rather than the rectus abdominis.

We observed similar inspiratory muscle activation patterns for EMG_{di} , EMG_{scm} and EMG_{para} between older and younger men at absolute and relative \dot{V}_E . Our finding contrasts Molgat-Seon et al. (2018b) who reported that EMG_{scm} was higher at absolute \dot{V}_E and EMG_{di} higher at absolute and relative \dot{V}_E , in older versus younger participants (men and women). It is possible that our results differ because women were not successfully recruited in our study, particularly given the sex differences in lung function and structure.

4.4 | Perceptual responses

We observed no differences between older and younger men for leg discomfort at absolute and relative \dot{V}_E . Older men had higher dyspnea ratings than younger men at absolute \dot{V}_E . However, at relative \dot{V}_E there were no dyspnea differences between the older and younger men. This agrees with other results reporting higher dyspnea ratings during submaximal exercise in older men compared to their younger counterparts (Ofir et al., 2008), but conflicts with others that found no difference between young and old men (Faisal et al., 2015). There were several differences in self-reported dyspnea descriptors between the age groups. Younger men were more likely to indicate that they were experiencing shallow breathing and expiratory difficulty than older men immediately following exercise. This is despite a lack of differences in their relative and absolute tidal volume or breathing frequency. Ofir et al. (2008) found that shallow breathing was reported by a greater proportion of older men, while Faisal et al. (2015) found no differences in dyspnea descriptors between younger and older adults. Our younger participants' perceptual results thus appear to contradict their physiological data highlighting the dichotomy between the biological and psychological influences of dyspnea perception.

An objective analysis of physiological data thus indicates that younger men's experience of dyspnea should be similar to that of the older men, yet their subjective experience was more unpleasant. The intensity of dyspnea during exercise is reduced by a recent experience of heightened dyspnea (Chang et al., 2023) and that exercising in room air, but believing that air to be hypoxic, results in greater dyspnea (Kipp et al., 2022). These results indicate that desensitization and expectation may play a role in the experience of dyspnea during exercise. It is also possible that desensitization and expectation could impact other aspects of the experience of dyspnea. We therefore suggest that differences in reporting of shallow breathing between our younger and older participants could be due to older participants being more habituated to the feeling of dyspnea and leg discomfort during exercise than our less active younger participants (Sucec et al., 2019). Alternatively, it could indicate that younger and older men undertaking exercise at similar intensities have different subjective experiences of exercise intensity. However, additional studies would be required to confirm either explanation.

Given that we observed higher expiratory muscle pressure generation during exercise in older men, it is natural to consider whether abdominal muscle fatigue may occur. In younger men, abdominal muscle fatigue can be induced by high-intensity constant load exercise (Hardy et al., 2021; Taylor et al., 2006; Taylor et al., 2013; Taylor & Romer, 2008; Verges et al., 2006), but whether this also occurs in healthy older men is unknown. Abdominal muscle fatigue could have implications for dyspnea, exercise tolerance, and diaphragm fatigue.

4.5 | Limitations

The main limitation to this study is its sample size which raises the possibility of type II errors. This occurred due to challenges in recruitment, primarily due to the time commitments, perceived invasiveness of testing, and restrictions on human testing due to COVID-19. Despite our original study design including men and women, only physically active older men were recruited. Though several older women expressed an interest to participate, most were excluded based on health risk factors during screening, and of those accepted into the study ($n=4$) a significant portion withdrew following falls at home ($n=2$) or after citing personal reasons ($n=1$). Younger women perceived the esophageal catheter as too invasive, resulting in very limited recruitment. Sample size was determined from power calculations based upon P_{ga} swings during exercise, potentially resulting in underpowered comparisons of measurements such as dyspnea and respiratory neuromuscular activation. Other limitations

include the potential for surface EMG recordings to be influenced by "cross talk" from nearby muscle groups. For instance, Abraham et al. (2002) reported that fine wire EMG_{ra} versus surface EMG_{ra} contained lower signal to noise, less artifacts, and recorded a wider range of motor unit sizes. Surface EMG_{para} has also been reported to be a poor surrogate for actual muscle activity compared to fine wire EMG_{para} (Tagliabue et al., 2021), although this conclusion has been questioned (Suh et al., 2021). In addition, EMG_{para} increases when participants cycle with their hands on the ergometer handlebars, likely due to coactivation of the pectoralis muscles (Ramsook et al., 2017). Similarly, EMG_{scm} may be influenced by activation of the scalenes (Mitchell et al., 2018). As the activation of confounding muscle groups cannot be eliminated, we instead minimized their influence by adhering to procedures designed to optimize signal quality. This included skin preparation, participant positioning during exercise, and instructions to minimize excess motion. Future research may benefit from the utilization of fine wire EMG due to its potential to limit cross talk. Surface EMG recordings were normalized against a maximal level of activity achieved during a maximal inspiratory or expiratory pressure, forced vital, or inspiratory capacity maneuver, to limit their susceptibility to interindividual artifacts. EMG_{di} was only measured from the crural, and not the costal, diaphragm during exercise. Thus, our EMG_{di} data does not represent the activation patterns of the diaphragm's entire musculature. Finally, P_{ga} may not fully capture the work completed by the contributions of the four muscle groups of the abdominal wall and is thus only an estimate of abdominal wall musculature's workload.

5 | CONCLUSION

We compared respiratory muscle pressure generation and inspiratory and expiratory neuromuscular recruitment patterns between younger and older men during exercise, alongside descriptors of dyspnea. Older men demonstrated greater expiratory muscle pressure generation during exercise compared to younger men and similar respiratory neuromuscular activation patterns for the diaphragm, sternocleidomastoids, parasternal intercostals and rectus abdominis. We also observed that younger men were more likely to indicate that they experienced shallow breathing than older men during exercise. Elevated expiratory muscle pressures may be due to compensatory mechanisms designed to offset increases in airway resistance due to aging. Future research is required to determine whether older men develop abdominal muscle fatigue in response to constant load high intensity exercise.

AUTHOR CONTRIBUTIONS

William MacAskill: conceptualization, methodology, software, validation, formal analysis, investigation, data curation, writing—original draft, visualization. **Ben Hoffman:** conceptualization, methodology, validation, formal analysis, writing—review and editing. **Michael A. Johnson:** formal analysis, methodology, writing—review and editing. **Graham R. Sharpe:** methodology, writing—review and editing. **Joshua Rands:** investigation, data curation, writing—review and editing. **Shoena E. Wotherspoon:** writing—review and editing formal analysis. **Yaroslav Gevorkov:** methodology, formal analysis, writing—review and editing. **Tracy L Kolbe-Alexander:** conceptualization, methodology, formal analysis, investigation, resources writing—review and editing supervision (equal). **Dean E. Mills:** conceptualization, methodology, software, validation, formal analysis, resources data curation, writing—review and editing, visualization, supervision (equal).

ACKNOWLEDGMENTS

The authors would like to acknowledge the assistance of Dr Johannes Moolman in screening potential participants for inclusion in this study. The authors would like to thank all participants for their time and dedication to this study. Open access publishing facilitated by University of Southern Queensland, as part of the Wiley - University of Southern Queensland agreement via the Council of Australian University Librarians.

FUNDING INFORMATION

University of Southern Queensland. Australian Government Research Training Program Scholarship.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

ETHICS STATEMENT

The study was approved by the University of Southern Queensland's Human Research Ethics Committee and all procedures conformed to the standards set by the Declaration of Helsinki, except for registration in a database (H17REA261).

ORCID

William MacAskill  <https://orcid.org/0000-0003-3979-5368>

REFERENCES

Abe, T., Sakamaki, M., Yasuda, T., Bembem, M. G., Kondo, M., Kawakami, Y., & Fukunaga, T. (2011). Age-related, site-specific muscle loss in 1507 Japanese men and women aged 20 to 95 years. *Journal of Sports Science and Medicine*, *10*, 145–150.

Abraham, K. A., Feingold, H., Fuller, D. D., Jenkins, M., Mateika, J. H., & Fregosi, R. F. (2002). Respiratory-related activation of human abdominal muscles during exercise. *Journal of Physiology*, *541*, 653–663.

Aliverti, A., Cala, S., Duranti, R., Ferrigno, G., Kenyon, C., Pedotti, A., Scano, G., Sliwinski, P., Macklem, P. T., & Yan, S. (1997). Human respiratory muscle actions and control during exercise. *Journal of Applied Physiology*, *1985*(83), 1256–1269.

AMERICAN THORACIC SOCIETY. (1999). Dyspnea. Mechanisms, assessment, and management: A consensus statement. *American Journal of Respiratory and Critical Care Medicine*, *159*, 321–340.

Baydur, A., Behrakis, P. K., Zin, W. A., Jaeger, M., & Milic-Emili, J. (1982). A simple method for assessing the validity of the esophageal balloon technique. *The American Review of Respiratory Disease*, *126*, 788–791.

Borg, G. A. (1982). Psychophysical bases of perceived exertion. *Medicine & Science in Sports & Exercise*, *14*, 377–381.

Chang, J. C., Angus, S. A., Doherty, C. J., Thompson, B. P., Mann, L. M., Molgat-Seon, Y., & Dominelli, P. B. (2023). Perception of exercise-induced dyspnea after experimentally induced breathing discomfort. *Applied Physiology, Nutrition, and Metabolism*, *48*, 514–525.

Cory, J. M., Schaeffer, M. R., Wilkie, S. S., Ramsook, A. H., Puyat, J. H., Arbour, B., Basran, R., Lam, M., Les, C., & Macdonald, B. (2015). Sex differences in the intensity and qualitative dimensions of exertional dyspnea in physically active young adults. *Journal of Applied Physiology*, *1985*(119), 998–1006.

Dacha, S., Janssens, L., Rodrigues, A., Louvaris, Z., Janssens, L., Gosselink, R., & Langer, D. (2019). Comparison between manual and (semi-) automated analyses of esophageal diaphragm electromyography during endurance cycling in patients with COPD. *Frontiers in Physiology*, *10*, 885.

DE Troyer, A., & Boriak, A. M. (2011). Mechanics of the respiratory muscles. *Comprehensive Physiology*, *1*, 1273–1300.

Elliott, J. E., Greising, S. M., Mantilla, C. B., & Sieck, G. C. (2016). Functional impact of sarcopenia in respiratory muscles. *Respiratory Physiology & Neurobiology*, *226*, 137–146.

Erail, S., Bostanci, Ö., & Polat, A. V. (2022). Ultrasound assessment of diaphragm thickness in athletes. *International Journal of Morphology*, *40*, 376–383.

Estenne, M., Yernault, J. C., & Troyer, D. E. (1985). Rib cage and diaphragm-abdomen compliance in humans: Effects of age and posture. *Journal of Applied Physiology*, *1985*(59), 1842–1848.

Faisal, A., Webb, K. A., Guenette, J. A., Jensen, D., Neder, J. A., O'donnell, D. E., & Network, C. R. R. (2015). Effect of age-related ventilatory inefficiency on respiratory sensation during exercise. *Respiratory Physiology & Neurobiology*, *205*, 129–139.

Fogarty, M. J., Omar, T. S., Zhan, W.-Z., Mantilla, C. B., & Sieck, G. C. (2018). Phrenic motor neuron loss in aged rats. *Journal of Neurophysiology*, *119*, 1852–1862.

Fukuchi, Y. (2009). The aging lung and chronic obstructive pulmonary disease: Similarity and difference. *Proceedings of the American Thoracic Society*, *6*, 570–572.

Fuller, D., Sullivan, J., & Fregosi, R. F. (1996). Expiratory muscle endurance performance after exhaustive submaximal exercise. *Journal of Applied Physiology*, *80*, 1495–1502.

Guenette, J. A., Chin, R. C., Cheng, S., Dominelli, P. B., Raghavan, N., Webb, K. A., Neder, J. A., & O'donnell, D. E. (2014). Mechanisms of exercise intolerance in global initiative for

- chronic obstructive lung disease grade 1 COPD. *The European Respiratory Journal*, 44, 1177–1187.
- Hardy, T. A., How, S. C., & Taylor, B. J. (2021). The effect of preexercise expiratory muscle loading on exercise tolerance in healthy men. *Medicine and Science in Sports and Exercise*, 53, 421–430.
- Holcombe, S., Ejima, S., & Wang, S. (2017). Calcification of costal cartilage in the adult rib cage. Proceedings of the 2017 international IRCOBI conference on the biomechanics of injury, Sep 13–15 2017 Antwerp. *Belgium*, 17–106, 737–744.
- Janssens, J.-P., Pache, J.-C., & Nicod, L. (1999). Physiological changes in respiratory function associated with ageing. *The European Respiratory Journal*, 13, 197–205.
- Jolley, C. J., Luo, Y. M., Steier, J., Rafferty, G. F., Polkey, M. I., & Moxham, J. (2015). Neural respiratory drive and breathlessness in COPD. *The European Respiratory Journal*, 45, 355–364.
- Kipp, S., Leahy, M. G., Hanna, J. A., & Sheel, A. W. (2022). Perceived exertion and dyspnea while cycling during a hypoxic and hyperoxic placebo. *Applied Physiology, Nutrition, and Metabolism*, 47, 1134–1142.
- Kubo, A. (1994). Changes in abdominal muscle strength with respect to aging. *Nihon Ronen Igakkai Zasshi*, 31, 525–531.
- Lalley, P. M. (2013). The aging respiratory system—Pulmonary structure, function and neural control. *Respiratory Physiology & Neurobiology*, 187, 199–210.
- Lansing, R. W., Gracely, R. H., & Banzett, R. B. (2009). The multiple dimensions of dyspnea: Review and hypotheses. *Respiratory Physiology & Neurobiology*, 167, 53–60.
- Lavolette, L., & Laveneziana, P. (2014). Dyspnoea: A multidimensional and multidisciplinary approach. *The European Respiratory Journal*, 43, 1750–1762.
- Macaskill, W., Hoffman, B., Johnson, M. A., Sharpe, G. R., & Mills, D. E. (2021). Pressure measurement characteristics of a microtransducer and balloon catheters. *Physiol Rep [Online]*, 9, 1–13. <https://doi.org/10.14814/phy2.14831>
- Meiners, S., Eickelberg, O., & Königshoff, M. (2015). Hallmarks of the ageing lung. *The European Respiratory Journal*, 45, 807–827.
- Miller, M. R., Hankinson, J., Brusasco, V., Burgos, F., Casaburi, R., Coates, A., Crapo, R., Enright, P., Van Der Grinten, C., & Gustafsson, P. (2005). Standardisation of spirometry. *Standardisation of Spirometry. European Respiratory Journal*, 26, 319–338.
- Mitchell, R. A., Schaeffer, M. R., Ramscook, A. H., Wilkie, S. S., & Guenette, J. A. (2018). Sex differences in respiratory muscle activation patterns during high-intensity exercise in healthy humans. *Respiratory Physiology & Neurobiology*, 247, 57–60.
- Mittman, C., Edelman, N. H., Norris, A. H., & Shock, N. W. (1965). Relationship between chest wall and pulmonary compliance and age. *Journal of Applied Physiology*, 1985(20), 1211–1216.
- Molgat-Seon, Y., Dominelli, P. B., Ramscook, A. H., Schaeffer, M. R., Molgat Sereacki, S., Foster, G. E., Romer, L. M., Road, J. D., Guenette, J. A., & Sheel, A. W. (2018). The effects of age and sex on mechanical ventilatory constraint and dyspnea during exercise in healthy humans. *Journal of Applied Physiology*, 1985(124), 1092–1106.
- Molgat-Seon, Y., Dominelli, P. B., Ramscook, A. H., Schaeffer, M. R., Romer, L. M., Road, J. D., Guenette, J. A., & Sheel, A. W. (2018). Effects of age and sex on inspiratory muscle activation patterns during exercise. *Medicine and Science in Sports and Exercise*, 50, 1882–1891.
- Molgat-Seon, Y., Ramscook, A. H., Peters, C. M., Schaeffer, M. R., Dominelli, P. B., Romer, L. M., Road, J. D., Guenette, J. A., & Sheel, A. W. (2019). Manipulation of mechanical ventilatory constraint during moderate intensity exercise does not influence dyspnoea in healthy older men and women. *The Journal of Physiology*, 597, 1383–1399.
- Ng, J., Kippers, V., & Richardson, C. (1998). Muscle fibre orientation of abdominal muscles and suggested surface EMG electrode positions. *Electromyography and Clinical Neurophysiology*, 38, 51–58.
- Ofir, D., Laveneziana, P., Webb, K. A., Lam, Y. M., & O'donnell, D. E. (2008). Sex differences in the perceived intensity of breathlessness during exercise with advancing age. *Journal of Applied Physiology*, 1985(104), 1583–1593.
- Ozemek, C., Whaley, M. H., Finch, W. H., & Kaminsky, L. A. (2017). Maximal heart rate declines linearly with age independent of cardiorespiratory fitness levels. *The European Respiratory Journal*, 17, 563–570.
- Parshall, M. B., Schwartzstein, R. M., Adams, L., Banzett, R. B., Manning, H. L., Bourbeau, J., Calverley, P. M., Gift, A. G., Harver, A., & Lareau, S. C. (2012). An official American Thoracic Society statement: Update on the mechanisms, assessment, and management of dyspnea. *American Journal of Respiratory and Critical Care Medicine*, 185, 435–452.
- Prakash, Y., & Sieck, G. C. (1998). Age-related remodeling of neuromuscular junctions on type-identified diaphragm fibers. *Muscle & Nerve*, 21, 887–895.
- Quanjer, P. H., Stanojevic, S., Cole, T. J., Baur, X., Hall, G. L., Culver, B. H., Enright, P. L., Hankinson, J. L., Ip, M. S., & Zheng, J. (2012). Multi-ethnic reference values for spirometry for the 3–95-yr age range: The global lung function 2012 equations. *The European Respiratory Journal*, 40, 1324–1343.
- Ramscook, A. H., Koo, R., Molgat-Seon, Y., Dominelli, P. B., Syed, N., Ryerson, C. J., Sheel, A. W., & Guenette, J. A. (2016). Diaphragm recruitment increases during a bout of targeted inspiratory muscle training. *Medicine and Science in Sports and Exercise*, 48, 1179–1186.
- Ramscook, A. H., Mitchell, R. A., Bell, T., Calli, S., Kennedy, C., Lehmann, J., Thompson, M., Puyat, J. H., & Guenette, J. A. (2017). Is parasternal intercostal EMG an accurate surrogate of respiratory neural drive and biomarker of dyspnea during cycle exercise testing? *Respiratory Physiology & Neurobiology*, 242, 40–44.
- Roman, M. A., Rossiter, H. B., & Casaburi, R. (2016). Exercise, ageing and the lung. *The European Respiratory Journal*, 48, 1471–1486.
- Romer, L. M., & Polkey, M. I. (2008). Exercise-induced respiratory muscle fatigue: Implications for performance. *Journal of Applied Physiology*, 1985(104), 879–888.
- Schaeffer, M. R., Mendonca, C. T., Levangie, M. C., Andersen, R. E., Taivassalo, T., & Jensen, D. (2014). Physiological mechanisms of sex differences in exertional dyspnoea: Role of neural respiratory motor drive. *Experimental Physiology*, 99, 427–441.
- Silva, C. G. D. S., Kaminsky, L. A., Arena, R., Christle, J. W., Araújo, C. G. S., Lima, R. M., Ashley, E. A., & Myers, J. (2018). A reference equation for maximal aerobic power for treadmill and cycle ergometer exercise testing: Analysis from the FRIEND registry. *European Journal of Preventive Cardiology*, 25, 742–750.
- Simon, P. M., Schwartzstein, R. M., Weiss, J. W., Fencel, V., Teghtsoonian, M., & Weinberger, S. (1990). Distinguishable types of dyspnea in patients with shortness of breath. *The American Review of Respiratory Disease*, 142, 1009–1014.
- Smith, J. R., Cross, T. J., Van Iterson, E. H., Johnson, B. D., & Olson, T. P. (2018). Resistive and elastic work of breathing in older and

- younger adults during exercise. *Journal of Applied Physiology*, 1985(125), 190–197.
- Sucec, J., Herzog, M., Van Den Bergh, O., Van Diest, I., & Von Leupoldt, A. (2019). The effects of repeated dyspnea exposure on response inhibition. *Frontiers in Physiology*, 10, 663.
- Suh, E.-S., D'cruz, R. F., Ramsay, M., Steier, J., Jolley, C. J., Reilly, C. C., Hart, N., Moxham, J., Rafferty, G., & Murphy, P. B. (2021). Parasternal electromyography as a surrogate measure of neural respiratory drive: Practical application and validity of surface and implanted fine wire methods. *Respiratory Physiology & Neurobiology*, 287, 103602.
- Tagliabue, G., Ji, M., Jagers, J. V. S., Lee, W., Dean, D., Zuege, D. J., Wilde, E. R., & Easton, P. A. (2021). Limitations of surface EMG estimate of parasternal intercostal to infer neural respiratory drive. *Respiratory Physiology & Neurobiology*, 285, 103572.
- Taylor, B. J., How, S. C., & Romer, L. M. (2006). Exercise-induced abdominal muscle fatigue in healthy humans. *Journal of Applied Physiology*, 1985(100), 1554–1562.
- Taylor, B. J., How, S. C., & Romer, L. M. (2013). Expiratory muscle fatigue does not regulate operating lung volumes during high-intensity exercise in healthy humans. *Journal of Applied Physiology*, 1985(114), 1569–1576.
- Taylor, B. J., & Romer, L. M. (2008). Effect of expiratory muscle fatigue on exercise tolerance and locomotor muscle fatigue in healthy humans. *Journal of Applied Physiology*, 1985(104), 1442–1451.
- Ticinesi, A., Meschi, T., Narici, M. V., Lauretani, F., & Maggio, M. (2017). Muscle ultrasound and sarcopenia in older individuals: A clinical perspective. *Journal of the American Medical Directors Association*, 18, 290–300.
- Turner, J. M., Mead, J., & Wohl, M. E. (1968). Elasticity of human lungs in relation to age. *Journal of Applied Physiology*, 1985(25), 664–671.
- Verges, S., Schulz, C., Perret, C., & Spengler, C. M. (2006). Impaired abdominal muscle contractility after high-intensity exhaustive exercise assessed by magnetic stimulation. *Muscle & Nerve*, 34, 423–430.
- Weavil, J. C., Thurston, T. S., Hureau, T. J., Gifford, J. R., Aminizadeh, S., Wan, H.-Y., Jenkinson, R. H., & Amann, M. (2022). Impact of aging on the work of breathing during exercise in healthy men. *Journal of Applied Physiology*, 132, 689–698.

How to cite this article: MacAskill, W., Hoffman, B., Johnson, M. A., Sharpe, G. R., Rands, J., Wotherspoon, S. E., Gevorkov, Y., Kolbe-Alexander, T. L., & Mills, D. E. (2023). The effects of age on dyspnea and respiratory mechanical and neural responses to exercise in healthy men. *Physiological Reports*, 11, e15794. <https://doi.org/10.14814/phy2.15794>